

Avast!® SC

DOC ID 585576

ACCEPTED FOR REGISTRATION
ONLY IN CONJUNCTION WITH
NEW YORK STATE SPECIFIC
SUPPLEMENTAL LABELING

1070.41

February 9, 2023

Classified for
“RESTRICTED USE”
in New York State
under 6NYCRR Part 326

New York State Department of
Environmental Conservation
Division of Materials Management
Pesticide Product Registration



FLURIDONE

GROUP

12

HERBICIDE

An herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, potable water sources, drainage canals and irrigation canals.

For use in New York State, comply with Section 24(C) Special Local Need labeling for Avast! SC, SLN NY-050003

Active Ingredient

Fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1*H*)-pyridinone 41.7%

Other Ingredients..... 58.3%

TOTAL 100.0%

Contains 4 pounds of fluridone per gallon.

Keep Out of Reach of Children CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

Refer to label booklet for additional Precautionary Information and Directions for Use including First Aid and Storage and Disposal.

Notice: Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse statements* in label booklet. If terms are unacceptable, return at once, unopened.**

EPA Reg. No. 67690-30
EPA Est. No. 067690-NC-001

FPL20221003
167002

SePRO Corporation 11550 N. Meridian St., Ste. 600, Carmel, IN 46032 U.S.A.

Aquatic Herbicide

Net contents 1 gallon (Non-refillable)

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053 .	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Gloves are required for the following application scenarios:

- Mixing/loading/applying with hand wand sprayer to ponds/lakes or static canals.
- Mixing/loading/applying with backpack sprayer to static canals.

ENGINEERING CONTROLS (AIRCRAFT)

Aircraft pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides 40 CFR 170.305.*

*Not for use in California.

ENVIRONMENTAL HAZARDS

Follow use directions carefully so as to minimize adverse effects on non-target organisms. Do not contaminate untreated water when disposing of equipment washwaters. Trees and shrubs growing in water treated with this product may occasionally develop chlorosis. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas.

Non-Target Organisms Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Shake well before using.

PRODUCT INFORMATION

This product is a selective systemic aquatic herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, drainage canals and irrigation canals. This product is absorbed from water by plant shoots and from hydrosol by the roots of aquatic vascular plants. It is important to maintain the specified concentration of this product in contact with the target plants for a minimum of 45 days. Rapid water movement or any condition that results in rapid dilution of this product in treated water will reduce its effectiveness.

In susceptible plants, this product inhibits the formation of carotene. In the absence of carotene, chlorophyll is rapidly degraded by sunlight. Herbicidal symptoms of this product appear in seven to ten days and appear as white (chlorotic) or pink growing points. Under optimum conditions, 30 to 90 days are required before the desired level of aquatic weed management is achieved with this product. Species susceptibility to this product may vary, depending on time of year, stage of growth, and water movement. For best results, apply this product prior to initiation of weed growth or when weeds begin active growth. Mature target plants may require an application rate at the higher end of the specified rate range and may take longer to control.

This product is not corrosive to application equipment.

This label provides recommendations on the use of a laboratory analysis for the active ingredient. SePRO Corporation recommends the use of high-performance liquid chromatography (HPLC) for the determination of fluridone concentrations in water. It is recommended to contact SePRO Corporation for the incorporation of this test, known as a FasTEST, in a treatment program. FasTEST is referenced in this label as the preferred method for the rapid determination of the active ingredient in water. Other proven chemical analyses for the active ingredient may also be used.

Application rates are provided in fluid ounces or quarts of this product to achieve a desired concentration of the active ingredient in parts per billion (ppb). **The maximum application rate or sum of all application rates is 90 ppb in ponds and 150 ppb in lakes and reservoirs per annual growth cycle.** This maximum concentration is the amount of product calculated as the target application rate, NOT determined by testing the residues of the active ingredient in the treated water.

Weed Resistance Management

For resistance management, this product is a Group 12 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 12 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 12 herbicides within a growing season or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers mechanical control methods, cultural (e.g., timing to favor the desirable plants and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular

weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your sales representative, pest control advisors, or local extension specialist for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of plants and weed biotypes.

Use Restrictions

- **Obtain required permits:** Permits may be required by state or local agencies. Consult with appropriate State or local water authorities before applying this product.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- **Hydroponic Farming:** Do not use water treated with this product for hydroponic farming unless a FasTEST has been run and confirmed that concentrations are less than 1 ppb.
- **Greenhouse and Nursery Plants:** Consult with SePRO Corporation for site-specific recommendations prior to any use of this product treated water for irrigating greenhouse or nursery plants. Without site-specific guidance from SePRO, do not use this product treated water for irrigating greenhouse or nursery plants unless a FasTEST has been run and confirmed that concentrations are less than 1 ppb..
- **Water Use Restrictions Following Applications of This Product (Days)**

Application Rate	Drinking ¹	Fishing	Swimming	Livestock/Pet Consumption	Irrigation ²
Maximum Rate (150 ppb) or less	0	0	0	0	See irrigation instructions below

¹ Note below, under *Potable Water Intakes*, the information for application of this product within ¼ mile (1,320 feet) of a functional potable water intake.

² Note below, under *Irrigation*, specific time frames or fluridone residues that provide the widest margin of safety for irrigating with water treated with this product.

- **Potable Water Intakes:** In lakes and reservoirs or other sources of potable water, do not apply this product at application rates greater than 20 ppb within ¼ mile (1,320 feet) of any functioning potable water intake. At application rates of 6 to 20 ppb, this product may be applied where functioning potable water intakes are present. **NOTE: Existing potable water intakes that are no longer in use, such as those that have been replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water intakes.**
- Aircraft pilots must use an enclosed cab that meets the definition listed in the WPS for agricultural pesticides 40 CFR 170.305.*

*Not for use in California.

Use Precautions

- **Irrigation:** Irrigation with water treated with this product may result in injury to the irrigated vegetation. Inform those who irrigate from areas treated with this product of the irrigation time frames or FasTEST requirements presented in the table below. Follow the following time frames and assay directions to reduce the potential for injury to vegetation irrigated with water treated with this product. There is a greater potential for crop injury when water treated with this product is applied to crops grown in low organic and sandy soils.

WAITING PERIODS BEFORE IRRIGATING WITH WATER TREATED WITH THIS PRODUCT			
Application Site	Days After Application		
	Established Tree Crops	Established Row Crops/ Turf/Plants	Newly Seeded Crops/Seedbeds or Areas to be Planted Including Overseeded Golf Course Greens
Ponds and Static Canals [†]	7	30	Assay required
Canals	7	14	Assay required
Lakes and Reservoirs ^{††}	7	14	Assay required

[†] For purposes of this product labeling, a pond is defined as a body of water 10 acres or less in size. A lake or reservoir is greater than 10 acres.

^{††} In lakes and reservoirs where one-half or greater of the body of water is treated, use the pond and static canal irrigation precautions.

Where the use of this product treated water is desired for irrigating crops prior to the time frames established above, the use of FastEST is recommended to measure the concentration in the treated water. Where FastEST has determined that concentrations are less than 10 parts per billion (ppb), there are no irrigation precautions for irrigating established tree crops, established row crops or turf. **For tobacco, tomatoes, peppers or other plants within the Solanaceae family and for newly seeded grasses, such as overseeded golf course greens, do not use this product treated water if concentrations are greater than 5 ppb. Furthermore, when rotating crops, do not plant members of the Solanaceae family in land that has been previously irrigated with fluridone concentrations in excess of 5 ppb. It is recommended that an aquatic specialist be consulted prior to commencing irrigation of these sites.**

WEED CONTROL INFORMATION

This product selectivity is dependent upon dosage, time of year, stage of growth, method of application and water movement. The following categories, Controlled, Partially Controlled and Not Controlled, are provided to describe expected efficacy under ideal treatment conditions, using higher to maximum application rates. Use of lower rates will increase selectivity of some species listed as Controlled or Partially Controlled. Additional aquatic plants may be controlled, partially controlled or tolerant to this product. Consult an aquatic specialist prior to application to determine a plant's susceptibility to this product.

Vascular Aquatic Plants Controlled

Floating Plants

Duckweed, Common (*Lemna minor*)

Emersed Plants

Spatterdock (*Nuphar luteum*)

Waterlily (*Nymphaea* spp.)

Submersed Plants

Bladderwort (*Utricularia* spp.)

Coontail, Common (*Ceratophyllum demersum*)

Egeria; Brazilian Elodea (*Egeria densa*)

Elodea, Common (*Elodea canadensis*)

Fanwort; Cabomba (*Cabomba caroliniana*)

Hydrilla (*Hydrilla verticillata*)

Naiad (*Najas* spp.)

Pondweed (*Potamogeton* spp.), except Illinois Pondweed

Watermilfoil (*Myriophyllum* spp.), except Variable-Leaf Milfoil

Shoreline Grasses

Paragrass (*Urochloa mutica*)

Vascular Aquatic Plants Partially Controlled**Floating Plants**

Salvinia (*Salvinia* spp.)

Watermeal, Common (*Wolffia columbiana*) †

† Partial control only with this product applied at the maximum labeled rate.

Emerged Plants

Alligatorweed (*Alternanthera philoxeroides*)

Cattail (*Typha* spp.)

Lotus, American (*Nelumbo lutea*)

Parrotfeather (*Myriophyllum aquaticum*)

Smartweed (*Polygonum* spp.)

Spikerush (*Eleocharis* spp.)

Waterprimrose, Creeping (*Ludwigia peploides*)

Waterpurslane (*Ludwigia palustris*)

Watershield (*Brasenia schreberi*)

Submersed Plants

Limnophila (*Limnophila sessiliflora*)

Pondweed, Illinois (*Potamogeton illinoensis*)

Tapegrass; American Eelgrass (*Vallisneria americana*)

Watermilfoil, Variable-Leaf (*Myriophyllum heterophyllum*)

Shoreline Grasses

Barnyardgrass (*Echinochloa crusgalli*)

Canarygrass, Reed (*Philaris arundinaceae*)

Cutgrass, Giant (*Zizaniopsis miliacea*)

Torpedograss (*Panicum repens*)

Watergrass, Southern (*Hydrochloa caroliniensis*)

Vascular Aquatic Plants Not Controlled**Floating Plants**

Water Lettuce (*Pistia stratiotes*)

Emerged Plants

Arrowhead (*Sagittaria* spp.)

Bacopa (*Bacopa* spp.)

Big Floatingheart; Banana Lily (*Nymphoides aquatica*)

Bulrush (*Scirpus* spp.)

Frogbit, American (*Limnobium spongia*)

Pickerelweed; Lanceleaf (*Pontederia* spp.)

Rush (*Juncus* spp.)

Waterhyacinth, Floating (*Eichornia crassipes*)

Water Pennywort (*Hydrocotyle umbellata*)

Shoreline Grasses

Maidencane (*Panicum hemitomom*)

NOTE: Algae (*Chara*, *Nitella* and filamentous species) are not controlled by this product.

MIXING AND APPLICATION DIRECTIONS

The aquatic plants present in the treatment site should be identified prior to application to determine their susceptibility to this product. It is also important to determine the area (acres) to be treated and the average depth in order to select the proper application rate. Do not exceed the maximum labeled rate for a given treatment site per annual growth cycle.

Shake this product well before using. Add the specified amount of this product to water in the spray tank during the filling operation. Agitate while filling and during spraying. Surface or subsurface application of the spray can be made with conventional spray equipment. This product can also be applied near the surface of the hydrosoil using weighted trailing hoses. A spray volume of 5 to 100 gallons per acre may be used. This product may also be diluted with water and the concentrated mix metered into the pumping system.

Tank Mix Directions

This product may be tank mixed with other aquatic herbicides and algaecides to enhance efficacy and plant selectivity. Refer to the label of the companion herbicide or algaecide for use directions, precautions and restrictions.

Application to Ponds

This product may be applied to the entire surface area of a pond. For single applications, rates may be selected to provide 45 to 90 ppb in the treated water. Use the higher rate within the rate range where there is a dense weed mass, when treating more difficult to control species, and for ponds that are less than 5 acres in size with an average depth of less than 4 feet. Application rates necessary to obtain these active ingredient concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the section of this label entitled *Application Rate Calculation - Ponds, Lakes and Reservoirs*. Split or multiple applications may be used where dilution of treated water is anticipated; however, the sum of all applications must not exceed a total of 90 ppb per annual growth cycle.

Average Water Depth of Treatment Site (feet)	Quarts of This Product per Treated Surface Acre			Fluid Ounces of This Product per Treated Surface Acre		
	45 ppb	to	90 ppb	45 ppb	to	90 ppb
1	0.12		0.24	3.8		7.7
2	0.24		0.49	7.7		15.7
3	0.37		0.73	11.8		23.4
4	0.49		0.98	15.7		31.4
5	0.61		1.22	19.5		39.0
6	0.73		1.46	23.4		46.7
7	0.85		1.70	27.2		54.4
8	0.98		1.95	31.4		62.4
9	1.10		2.19	35.2		70.1
10	1.22		2.44	39.0		78.1

Application to Lakes and Reservoirs

The following treatments may be used for treating both whole lakes or reservoirs and partial areas of lakes or reservoirs (bays, etc.). For best results in treating partial lakes and reservoirs, this product treatment areas should be a minimum of 5 acres in size. Treatment of areas smaller than 5 acres or treatment of narrow strips, such as boat lanes or shorelines, may not produce satisfactory results due to dilution by untreated water. Rate ranges are provided as a guide to include a wide range of environmental factors, such as target species, plant susceptibility, selectivity and other aquatic plant management objectives. Application rates and methods should be selected to meet the specific lake/reservoir aquatic plant management goals.

Whole Lake or Reservoir Treatments (Limited or No Water Discharge)

Single Application to Whole Lakes or Reservoirs

Where single applications to whole lakes or reservoirs are desired, this product may be applied at an application rate of 10 to 90 ppb. Application rates necessary to obtain these concentrations in treated water are shown in the following table. For additional application rate calculations, refer to the section of this label entitled *Application Rate Calculation - Ponds, Lakes and Reservoirs*. Choose an application rate from the table below to meet the aquatic plant management objective. **Where greater plant selectivity is desired, such as when controlling Eurasian watermilfoil and curlyleaf pondweed, an application rate lower in the rate range may be chosen.** For other plant species, an aquatic specialist should be contacted to determine when to choose application rates lower in the rate range to meet specific plant management goals. Use the higher rate within the rate range where there is a dense weed mass or when treating more difficult to control plant species. Retreatments may be required to control more difficult to control species or in the event of a heavy rainfall event where dilution of the treatment concentration has occurred. In these cases, a second application or more may be required; however, the sum of all applications must not exceed 150 ppb per annual growth cycle. Refer to the following section, *Split or Multiple Applications to Whole Lakes or Reservoirs*, for guidelines and maximum rate allowed.

Average Water Depth of Treatment Site (feet)	Quarts of This Product per Treated Surface Acre			Fluid Ounces of This Product per Treated Surface Acre		
	10 ppb	to	90 ppb	10 ppb	to	90 ppb
1	0.03		0.24	1.0		7.7
2	0.05		0.49	1.6		15.7
3	0.08		0.73	2.6		23.4
4	0.11		0.98	3.2		31.4
5	0.14		1.22	4.5		39.0
6	0.16		1.46	5.1		46.7
7	0.19		1.70	6.1		54.4
8	0.22		1.95	7.0		62.4
9	0.24		2.19	7.6		70.1
10	0.27		2.44	8.6		78.1
11	0.30		2.68	9.6		86.0
12	0.32		2.93	10.2		93.8
13	0.35		3.17	11.2		101.4
14	0.38		3.42	12.1		109.4
15	0.41		3.66	13.1		117.1
16	0.43		3.90	13.8		124.8
17	0.46		4.15	14.7		132.2
18	0.49		4.39	15.7		140.5
19	0.51		4.63	16.3		148.2
20	0.54		4.88	17.3		156.2

Split or Multiple Applications to Whole Lakes or Reservoirs

To meet certain plant management objectives, split or multiple applications may be desired in making whole lake treatments. Split or multiple application programs are desirable when the objective is to use the minimum effective dose and, through the use of a water analysis, e.g. FasTEST or other appropriate means of analysis, add additional product to maintain this lower dose for sufficient time to ensure efficacy and enhance selectivity. Water may be treated with an initial application of 6 to 50 ppb. Additional split applications should be made to maintain a sufficient concentration for a minimum of 45 days. **In controlling Eurasian watermilfoil and curlyleaf pondweed and where greater plant selectivity is desired, an application rate lower in the rate range may be chosen.** For other plant species, an aquatic specialist should be contacted to determine when to choose application rates lower in the rate range to meet specific plant management goals. When utilizing split or multiple applications of this product, the utilization of FasTEST is strongly recommended to determine the actual concentration in the water over time. For split or multiple applications, the sum of all applications must not exceed 150 ppb per annual growth cycle.

NOTE: In treating lakes or reservoirs that contain functional potable water intakes and the application requires treating within ¼ mile of a potable water intake, no single application can exceed 20 ppb. Additionally, the sum of all applications must not exceed 150 ppb per annual growth cycle.

Partial Lake or Reservoir Treatments

Where dilution of this product with untreated water is anticipated, such as in partial lake or reservoir treatments, split or multiple applications may be used to extend the contact time with the target plants. The application rate and use frequency of this product in a partial lake is highly dependent upon the treatment area. An application rate at the higher end of the specified rate range may be required and frequency of applications will vary depending upon the potential for untreated water to dilute the this product concentration in the treatment area. Use a rate at the higher end of the rate range where greater dilution with untreated water is anticipated.

Treatment Areas Greater than ¼ Mile from a Functioning Potable Water Intake

For single applications, this product may be applied at application rates from 30 to 150 ppb. Split or multiple applications may be made; however, the sum of all applications must not exceed 150 ppb per annual growth cycle. Split applications may be made to maintain a sufficient concentration in the target area for a period of 45 days or longer. The use of FasTEST, or other appropriate means of analysis, is recommended to maintain the desired concentration in the target area over time.

Treatment Areas Within ¼ Mile of a Functioning Potable Water Intake

In treatment areas that are within ¼ mile of a potable water intake, no single application can exceed 20 ppb. When utilizing split or multiple applications of this product for sites that contain a potable water intake, FasTEST or other appropriate means of analysis is required to determine the actual concentration in the water. Additionally, the sum of all applications must not exceed 150 ppb per annual growth cycle.

Application Rate Calculation — Ponds, Lakes and Reservoirs

The amount of this product to be applied to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

Quarts of this product required per treated surface acre = Average water depth of treatment site (feet) x Desired ppb concentration of active ingredient x 0.0027

For example, the quarts per acre of this product required to provide a concentration of 25 ppb of active ingredient in water with an average depth of 5 feet is calculated as follows:

$$5 \times 25 \times 0.0027 = 0.33 \text{ quarts per treated surface acre}$$

When measuring quantities of this product, quarts may be converted to fluid ounces by multiplying quarts to be measured by 32. For example, 0.33 quarts x 32 = 10.5 fluid ounces.

NOTE: Calculated rates may not exceed the maximum allowable rate in quarts per treated surface acre for the water depth listed in the application rate table for the site to be treated.

Application to Drainage Canals and Irrigation Canals

Static Canals

In static drainage and irrigation canals, apply this product at the rate of 1 to 2 quarts per treated surface acre.

Moving Water Canals

The performance of this product will be enhanced by restricting or reducing water flow. In slow moving bodies of water, use an application technique that maintains a concentration of 15 to 40 ppb in the target area for a minimum of 45 days. This product can be applied by split or multiple broadcast applications or by metering in the product to provide a uniform concentration of the herbicide based upon the flow pattern. The use of FastEST or other appropriate means of analysis is recommended to maintain the desired concentration in the target area over time.

Static or Moving Water Canals Containing a Functioning Potable Water Intake

In treating a static or moving water canal that contains a functioning potable water intake, DO NOT apply this product at application rates greater than 20 ppb within ¼ mile (1320 feet) of any functioning potable water intake. Applications of less than 20 ppb may be applied within ¼ mile from a functioning potable water intake; however, if applications of this product are made within ¼ mile from a functioning potable water intake, the FastEST or other appropriate means of analysis must be utilized to demonstrate that concentrations do not exceed 150 ppb at the potable water intake.

Application Rate Calculation – Moving Water Drainage Canals and Irrigation Canals

The amount of this product to be applied through a metering system to provide the desired ppb concentration of active ingredient in treated water may be calculated as follows:

1. Average flow rate (feet per second) x Average canal width (feet) x Average water depth (feet) x 0.9 = Cubic feet per second (CFS)
2. CFS x 1.98 = acre-feet per day (water movement)
3. Acre-feet per day x desired ppb x 0.0027 = Quarts of this product required per day

SPRAY DRIFT ADVISORIES

The applicator is responsible for avoiding off-site spray drift. Be aware of nearby non-target sites and environmental conditions.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.*
* Not for use in California.

Boom Height – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

Release Height - Aircraft

Higher release heights increase the potential for spray drift.*

*Not for use in California.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage:

Store in original container only. Do not store near feed or foodstuffs. Keep from freezing.

Pesticide Disposal:

Wastes resulting from use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity >5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for

continued

STORAGE AND DISPOSAL *continued*

later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.


Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

Warranty Disclaimer: SePRO Corporation warrants that this product conforms to the chemical description on the product label. Testing and research have also determined that this product is reasonably fit for the uses described on the product label. To the extent consistent with applicable law, SePRO Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty and any such warranties are expressly disclaimed.

Misuse: Federal law prohibits the use of this product in a manner inconsistent with its label directions. To the extent consistent with applicable law, the buyer assumes responsibility for any adverse consequences if this product is not used according to its label directions. In no case shall SePRO Corporation be liable for any losses or damages resulting from the use, handling or application of this product in a manner inconsistent with its label.

For additional important labeling information regarding SePRO Corporation's Terms and Conditions of Use, Inherent Risks of Use and Limitation of Remedies, please visit <http://seprolabels.com/terms> or scan the image below.



©Copyright 2022 SePRO Corporation

Avast! is a registered trademark of SePRO Corporation

Avast!® SC

1070.41



Active Ingredient

Fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1 <i>H</i>)-pyridinone.....	41.7%
Other Ingredients	58.3%
TOTAL	100.0%

Contains 4 pounds of fluridone per gallon of product.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC at 1-800-535-5053 .	

FLURIDONE	GROUP	12	HERBICIDE
------------------	--------------	-----------	------------------

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.
Pesticide Storage: Store in original container only. Do not store near feed or foodstuffs. Keep from freezing.
Pesticide Disposal: Wastes resulting from use of this product may be disposed of on-site or at an approved waste disposal facility.
Container Handling
Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.
See attached booklet for complete container disposal directions including triple rinsing and pressure rinsing instructions.

Refer to label booklet for additional Precautionary Information and Directions for Use.

NOTICE: Read the entire label before using. Use only according to label directions. **Before buying or using this product, read *Warranty Disclaimer* and *Misuse* statements inside label booklet. If terms are unacceptable, return at once, unopened.**

EPA Reg. No. 67690-30
 EPA Est. No. 067690-NC-001

167002
 FPL20221003

SePRO Corporation
 11550 N. Meridian Street, Suite 600 • Carmel, IN 46032, U.S.A.



Aquatic Herbicide

Net contents 1 gallon (Non-refillable)